

◆ Research paper

DOI: [10.5281/zenodo.14025324](https://doi.org/10.5281/zenodo.14025324)

Flora of Protected and Environmentally Sensitive Afforestation Area in Çankırı, Türkiye

Gamze TUTTU¹, Ebru GÜL^{1*}

¹Department of Forest Engineering, University of Çankırı, Türkiye

* Corresponding author Email: ebru@karatekin.edu.tr

Abstract: The objective of this study was to examine the flora of Çankırı Karatekin University Faculty of Forestry Prof. Dr. Abdulreşit Brohi Research and Application Forest and to provide insights that could contribute to the conservation of biodiversity in the region. This afforestation area is a 25-year-old protected area that is sensitive to desertification. Despite the maintenance and restoration efforts undertaken in the region, its combination of protected status and high sensitivity to desertification underscores this study's importance. Plant specimens were systematically collected from designated areas, and identification procedures were carried out following established botanical methods. Between April and August 2021, 205 plant specimens were gathered from the area. The identification process recorded 192 taxa representing 153 genera and 47 families. Among these taxa, one belongs to the *Pteridophyta* division, while 191 are classified under the *Spermatophyta* division. Notably, no natural Gymnosperm specimens were found in the area. Among the identified taxa, 170 are *Dicotyledoneae*, and 22 are *Monocotyledoneae* species. The families with the highest representation are as follows: *Asteraceae*, with 31 species; *Brassicaceae*, with 19 species; *Fabaceae*, comprising 17 species; *Lamiaceae*, consisting of 16 species; and *Poaceae*, which includes 15 species. The most abundant genera, represented by three species, are *Centaurea*, *Crepis*, *Tragopogon*, *Silene*, *Convolvulus*, *Astragalus*, *Onobrychis*, *Salvia*, *Teucrium* and *Veronica*. The study identified ten endemic taxa, resulting in an endemism ratio of 5.21%. These endemic taxa play a crucial role in conserving biodiversity in the region. In terms of phytogeographic regions, the taxa can be categorized as follows: Irano-Turanian (46 taxa, 23.96%), Euro-Siberian (9 taxa, 4.69%), Mediterranean (4 taxa,

2.08%), Eastern Mediterranean (3 taxa, 1.56%), with the remaining 130 taxa (67.71%) classified as unknown or multiregional. The research findings provide essential insights into regional conservation and sustainable management.

Keywords: afforestation, desertification, flora, protected area, semiarid

Introduction

Preserving semi-arid and desertification-sensitive areas is essential for ensuring the sustainability of ecosystems. Turkey is endowed with a remarkable biodiversity, which can be attributed to the country's diverse climate and varied geological structures. This situation is critically essential for Turkey's ecosystem functions and contribution to global biodiversity. A total of 11,707 taxa from the classes of ferns and seed plants with conduction bundles have been identified in Turkey, with one-third of these (3,649) being endemic (Güner et al., 2012). Turkey is recognized as one of the richest regions in the world concerning endemic plant diversity, with 34.4% of its total endemic species (Ekim, 2009; Tekeli et al., 2006; Demir, 2013). It is crucial to identify the habitats of primarily narrowly distributed endemic species to protect them and highlight their significance to our country (Tuttu, 2024).

Turkey's geographical location and climate characteristics place it within a semi-arid climate zone, exacerbating the inadequacy of water resources and the frequency of drought events across a considerable portion of the country. A considerable proportion of the total area of 51 million hectares is classified as arid and semi-arid regions, which suggests that the risk of drought will increase in the future (Edis et al., 2022; Serkendiz & Tatlı, 2023; Erişmiş, 2023; Edis, 2024). In arid and semi-arid regions, a reduction in soil properties and water retention capacity harms agricultural production. This, in turn, exacerbates the degradation of natural vegetation and the phenomenon of desertification, which are further compounded by climate change (Akbaş Tetik et al., 2023).

The semi-arid regions in Turkey are mainly prone to erosion, highlighting the essential need for afforestation efforts (Edis et al. 2023). In afforestation projects conducted in semi-arid regions, choosing species that can withstand harsh climatic conditions and enhance soil properties is crucial (Edis et al., 2021). Afforestation is a crucial tool in the fight against desertification (Kar, 2018), enhancing carbon sequestration (Lal, 2004), and promoting

sustainable development. However, it is essential to highlight that afforestation initiatives can significantly change ecosystem functions. These changes may include shading, microclimate, nutrient cycling, and water balance alterations, which can ultimately affect biodiversity (Elmarsdottir et al., 2008).

The sustainability of ecosystems is contingent upon the continued viability of semi-arid and desertification-sensitive areas. In such areas, the ability of plants to withstand challenges such as climate change, wind erosion, and drought is critical for the sustainability of ecosystems. In arid areas such as Central Anatolia, it has been observed that vegetation cover has been significantly degraded over many years due to human and animal pressure (Fırıncioğlu et al., 2007). It is, therefore, essential to identify suitable plant species to restore these ecosystems. It has been demonstrated that plant species in semi-arid regions demonstrate resilience to desertification processes and provide significant contributions to ecosystem services (Noy-Meir, 1973; Maestre et al., 2012). These species are notable for their capacity to utilize water efficiently and prevent soil erosion, which are critical factors in the success of afforestation efforts (López-Bermúdez et al., 1998). Plant species selection and adaptation are essential in semi-arid and desertification-sensitive afforestation areas.

The floristic composition of afforestation areas significantly influences biodiversity, determining the diversity and abundance of plant species. Urban and rural afforestation projects play a significant role in maintaining ecological balance. The objective of this study is to examine the plant species found in semi-arid and desertification-sensitive afforestation areas and to elucidate their impact on the ecosystem. Furthermore, evaluating these species' adaptation processes and their contribution to biodiversity will provide crucial data for developing effective ecosystem management and conservation policies in the region.

Material and Methods

Study area

The research and application forest at Çankırı Karatekin University (ÇAKÜ) Faculty of Forestry is 9.4 ha in size (Fig. 1). Since 1998, approximately 2,000 saplings of various species, including *Pinus nigra* J.F.Arnold subsp. *pallasiana* (Lamb.) Holmboe, *Cedrus libani* L., *Amygdalus communis* L., *Juniperus sabina* L., and *Robinia pseudoacacia* L., have been planted in the study

area. The climate of the study area can be described as follows. According to the meteorological data of Çankırı province (Anonymous, 2022), the average temperature is 11.1 °C, the average annual precipitation is 393.9 mm, the highest precipitation is in May with 53.6 mm, and the lowest precipitation is in September with 16.4 mm. By the classification proposed by Thornthwaite (1949), the area is characterized by a semi-arid climate.




Figure 1. Location of the study area (Gül et al., 2019)

Desertification tendency of the study area

The desertification risk and environmental sensitivity of the study area were previously determined by Gül et al. (2019). This section provides general information on the area's desertification tendency (for detailed information, see Gül et al., 2019). Gül et al. (2019) used the Desertification Indicator System for Mediterranean Europe (DIS4ME) system to assess the desertification tendency of the afforested area. As a result of the calculations, it was found that the risk of desertification in the study area was evaluated in the low-risk class with the lowest

1.69 and in the medium-risk class with the highest 5.25. The desertification risk was lower in areas where the vegetation cover and soil properties improved due to afforestation activities. These results show that afforestation activities in the area effectively prevent desertification. Similarly, the Environmentally Sensitive Area (ESA) sensitivity index showed that the area's sensitivity varied between 1.35 and 1.52. These values indicate that the area falls into critical and fragile sensitivity classes. Fragile areas become more vulnerable to desertification, especially if not protected.

Collection and identification of plant samples

The material under investigation comprises plant samples gathered systematically between April and August 2021. During the collection of plant specimens, particular attention was paid to flowers, fruits, seeds, and other relevant plant parts. At least two specimens of each individual were collected, prioritizing undamaged ones and distributing them evenly throughout the area. The specimens were carefully placed in plastic bags and transported to the laboratory, where they were subjected to drying processes using the methodologies established by Yaltırık and Efe (1996). The identified specimens are stored in the Department of Forest Botany laboratory at Çankırı Karatekin University.

The series "Flora of Turkey and the East Aegean Islands," which includes volumes 1-9 (Davis, 1965-1985) and a supplement (Davis et al., 1988), served as the primary source for plant identification. "Flora Europaea" (Tutin, 1964-1993) was also referenced. The key resources for plant identification were Volume 11 (Güner et al., 2000) and "Flora Europaea" (Tutin, 1964-1993). The identification process utilized a Leica S6T stereo research microscope. The English-Turkish Botany Guide (Baytop, 1998) and Plant Identification Terminology (Harris and Harris, 2001) were consulted to understand the meanings of morphological terms.

The endangered status of endemic species was determined using the "Red Book of Plants of Turkey" (Ekim et al., 2000), while current names and authors were sourced from the WFO Plant List and IPNI databases (<https://wfoplantlist.org/>; <http://www.ipni.org>). The "List of Plants of Turkey (Vascular Plants)" (Güner et al., 2012) was also referenced. Furthermore, the Virtual Herbariums of the Van Lake Basin (<http://www.vanherbaryum.yyu.edu.tr>) and the Berlin

Museum of Natural History (<http://ww2.bgbm.org/herbarium>) were used for identifying seed plants.

Results

The 205 plant specimens were collected from the area in 2021 between April and August (Table 1). As a result of the identification of plant specimens, 192 taxa belonging to 153 genera and 47 families have been determined. One belongs to the Pteridophyta, and 191 taxa to the Spermatophyta division. All plant specimens are in the Angiosperme section in the area. The 170 taxa are Dicotyledoneae, and 22 taxa are Monocotyledoneae. Regarding distribution according to phytogeographic regions, Irano-Turanian 46 (23.96%) and Euro-Siberian 9 (4.69%) and Mediterranean-E.Mediterranean 7 (3.64%) phytogeographic region. The remaining 130 (67.71%) taxa are distributed unknown or multiregional (Table 2).

Table 2. Distribution of plant species in the study area by phytogeographic region

Phytogeographic Region	Number of Taxa	Proportional Distribution (%)
Irano-Turanian element	46	23.96
Euro-Siberian element	9	4.69
Mediterranean element	7	3.64
Unknown or multiregional	130	67.71
Total	192	100

According to the Flora of Turkey and the Eastern Aegean Islands findings, the five most prevalent families in terms of species diversity are Asteraceae, Fabaceae, Lamiaceae, Brassicaceae, and Poaceae. In our study, the initial five families are identical. As Erik and Tarikahya (2004) state, Asteraceae, the most prominent family in Turkey, is represented by 1186 species, accounting for 13.1% of the country's flora. As illustrated in Table 3, the Asteraceae family is the most prevalent in the study area, comprising 31 taxa and accounting for 16.1% of the flora. Asteraceae is a cosmopolitan family exhibiting high ecological tolerance and the capacity for seed dispersal.

Table 3 The largest families

Family	Number of Taxa	Percentage of Total Number of Species
<i>Asteraceae</i>	31	16.1
<i>Brassicaceae</i>	19	9.9
<i>Fabaceae</i>	17	8.9
<i>Lamiaceae</i>	16	8.3
<i>Poaceae</i>	15	7.8
Other	94	49

Table 4 displays the taxon with the highest number of taxa in the research area. The largest genera include *Centaurea*, *Crepis*, *Tragopogon*, *Silene*, *Convolvulus*, *Astragalus*, *Onobrychis*, *Salvia*, *Teucrium*, and *Veronica*, with three examples from each. The largest genus of flora in Turkey, *Astragalus*, is distributed across the Irano-Turanian phytogeographic region and in habitats characterized by steppe. Despite its location in a transitional climate zone, the area is generally under the influence of the Iranian-Turanian Flora Region, which lends support to the conclusion that the genus *Astragalus* is the most prevalent (Erik and Tarikahya, 2004).

Table 4: The taxon that includes the highest number of taxa in the research area

Genus	Taxa	Percent %
<i>Astragalus</i>	3	1.56
<i>Centaurea</i>	3	1.56
<i>Convolvulus</i>	3	1.56
<i>Crepis</i>	3	1.56
<i>Onobrychis</i>	3	1.56
<i>Salvia</i>	3	1.56
<i>Silene</i>	3	1.56
<i>Teucrium</i>	3	1.56
<i>Tragopogon</i>	3	1.56
<i>Veronica</i>	3	1.56

Ten endemic plant species have been identified in the study area (Table 5), resulting in an endemism rate of 5.21%. These plants belong to the Iran-Turan phytogeographic region 5.21%.

Table 5: Endemic taxa in the study area and their associated threat categories (LR(lc): Lover risk, Least Concern)

	Family	Taxa	Threat Categories
1	Asteraceae	<i>Cirsium pseudocreticum</i> (P.H.Davis & Parris) Yıldız, Dirmenci & Arabacı	LR(lc)
2	Asteraceae	<i>Jurinea pontica</i> Hausskn. & Freyn ex Hausskn.	LR(lc)
3	Asteraceae	<i>Tragopogon aureus</i> Boiss.	LR(lc)
4	Boraginaceae	<i>Mattiastrum leptophyllum</i> Brand	LR(lc)
5	Boraginaceae	<i>Onosma isaurica</i> Boiss. & Heldr.	LR(lc)
6	Brassicaceae	<i>Odontarrhena pateri</i> (Nyár.) Španiel, Al-Shehbaz, D.A.German & Marhold	LR(lc)
7	Caryophyllaceae	<i>Gypsophila eriocalyx</i> Boiss.	LR(lc)
8	Fabaceae	<i>Onobrychis ornata</i> Desv.	LR(lc)
9	Lamiaceae	<i>Salvia absconditiflora</i> Greuter & Burdet	LR(lc)
10	Rubiaceae	<i>Cynanchica bornmuelleri</i> (Velen. ex Bornm.) P.Caputo & Del Guacchio	LR(lc)

Discussion and Conclusion

This study presents the findings of a comprehensive survey conducted to evaluate the plant species composition in a protected afforestation site located in a semi-arid region facing potential desertification. The results emphasize the distribution of different plant species within the ecosystem. A total of 192 plant taxa were identified, including 10 local endemic species. The flora study revealed that the most dominant species in the study area belong to the Asteraceae, Brassicaceae, Fabaceae, Lamiaceae, and Poaceae families. These species are closely related to the local climatic conditions and soil characteristics.

To ascertain the floristic composition, it is essential to compare studies conducted in neighboring areas to identify similarities and differences in the plant species present. The families with the highest number of identified taxa in the studies conducted in the study area and nearby areas were compared and are presented in Table 6. Upon analysis of this table, it becomes evident that the Asteraceae family has the highest number of taxa. When the other families ranked in all study areas are examined, it is observed that there are some differences. This can be attributed to various factors, including the size of the areas, spatial differences, variability in climate and soil characteristics, vegetation types, and differences in vegetation layers.

Table 6. Comparison of Families with the Highest Number of Taxa

Family	Neighboring Studies								
	1	2	3	4	5	6	7	8	9
	Taxa (%)	Taxa (%)	Taxa (%)	Taxa (%)	Taxa (%)	Taxa (%)	Taxa (%)	Taxa (%)	Taxa (%)
Asteraceae	31 (16.1)	45 (14.8)	52 (14.6)	75 (11.9)	51 (14.1)	66 (15.8)	35 (15.0)	33 (14.0)	35 (14.0)
Brassicaceae	19 (9.9)	11 (3.62)	21 (5.88)	44 (6.98)	35 (9.6)	21 (5.03)	16 (6.87)	13 (5.5)	8 (3.2)
Fabaceae	17 (8.9)	29 (9.54)	33 (9.24)	48 (7.61)	39 (10.8)	44 (11.5)	32 (13.7)	28 (11.9)	26 (10.4)
Lamiaceae	16 (8.3)	24 (7.9)	34 (9.52)	43 (6.82)	29 (8.0)	41 (9.83)	25 (10.7)	31 (13.2)	24 (9.6)
Poaceae	15 (7.8)	9 (2.96)	12 (3.36)	56 (8.88)	30 (8.3)	18 (4.31)	9 (3.87)	11 (4.7)	15 (6.0)
Other	94 (49)	186 (6.1.1)	205 (57.4)	364 (57.7)	181 (49.6)	227 (54.4)	116 (49.8)	119 (50.6)	144 (57.1)
Total	192 (100)	304 (100)	357 (100)	630 (100)	365 (100)	417 (100)	233 (100)	235 (100)	252 (100)

1 ÇAKÜ Abdülreşit Brohi Ağaçlandırma Sahası (Tuttu and Gül, 2024), **2** Research and Application Forest (Eldivan) (Tuttu and Ursavaş, 2022), **3** Çankırı-Korubaşı Hill and its surroundings (Tuttu and Akkemik, 2017), **4** Ilgaz Mountain National Park (Pehlivan, 2007), **5** Karliktepe (Çankırı) (Sağıroğlu and Duman, 2004), **6** Çankırı Yapraklı Forests (Mutlu, 2006), **7** Büyükyayla (Yapraklı) (Dörlarslan and Gül, 2015), **8** Küçükyayla (Yapraklı) (Dörlarslan and Gül, 2021), **9** Kabalı Dağı (Çerkeş) Florası (Erdoğan, Ketenoglu and Bingöl, 2008)

The distribution of the taxa, according to phytogeographic regions; Irano-Turanian 46 (23.96%), Euro-Siberian 9 (4.69%), Mediterranean 4 (2.08%), E.Mediterranean 3 (1.56%) and unknown or multiregional taxa are 130 (67.71%), (Tablo7). As illustrated in Table 7, the greatest concentration of Irano-Turanian elements is observed within the study area. Except for the flora of Ilgaz Mountain National Park, the proportion of Irano-Turanian elements was notably elevated across all studies. Ilgaz Mountain National Park is situated at a higher elevation in the northern region than other areas, so there is a greater prevalence of Euro-Siberian elements.

Table 7: Distribution of other studies conducted in and around the research area, classified according to phytogeographical regions.

Phytogeographic Regions	Neighboring Studies								
	1	2	3	4	5	6	7	8	9
Irano-Turanian element (%)	23,96	16,77	27,20	12,00	27,70	20,60	24,03	20,00	14,80
Euro-Siberian element (%)	4,69	14,14	8,10	33,00	4,10	15,50	12,87	11,50	10,40
Mediterranean element (%)	3,64	2,96	5,30	6,00	3,60	3,80	3,87	5,10	4,00
Unknown or multiregional (%)	67,71	60,53	59,40	49,00	64,60	60,10	59,23	63,40	70,80
Taxa	192	304	357	630	365	417	233	235	252

1 ÇAKÜ Abdülreşit Brohi Ağaçlandırma Sahası (Tuttu and Gül, 2024), **2** Research and Application Forest (Eldivan) (Tuttu and Ursavaş, 2022), **3** Çankırı-Korubaşı Hill and its surroundings (Tuttu and Akkemik, 2017), **4** Ilgaz Mountain National Park (Pehlivan, 2007), **5** Karlıktepe (Çankırı) (Sağıroğlu and Duman, 2004), **6** Çankırı Yapraklı Forests (Mutlu, 2006), **7** Büyükyayla (Yapraklı) (Dörlarslan and Gül, 2015), **8** Küçükyayla (Yapraklı) (Dörlarslan and Gül, 2021), **9** Kabalı Dağı (Çerkeş) Florası (Erdoğan, Ketenoglu and Bingöl, 2008)

In conclusion, this study provides a valuable resource for understanding semi-arid protected and environmentally sensitive afforestation areas' floristic composition and ecosystem dynamics. Flora studies are essential for understanding ecosystem functioning, conserving biodiversity, and ensuring environmental sustainability. These studies provide the necessary information to address environmental problems at both local and global scales and form a critical basis for maintaining ecosystem health.

Reference

- Akbaş Tetik, N., Gül, E., Dörlarslan, M. (2023). Evaluation of desertification tendency based on soil characteristics in Çankırı urban forest. Anatolian Journal of Forest Research, 9(2), 101-106.
- Anonymous. 2021. Çankırı Meteorology Bulletin, Republic of Turkey Ministry of Environment, Urbanization and Climate Change, General Directorate of Meteorology Records, Ankara.
- Baytop, A., 1998. İngilizce-Türkçe Botanik Kılavuzu, *İ.Ü.Basimevi, Eczacılık Fakültesi Yayın No: 70, İstanbul.*

- Davis, P.H., 1965-1985. Flora of Turkey and the East Aegean Islands, Vol. 1-9, *Edinburg University Press, Edinburgh*.
- Davis, P.H., Mill, R.R., Tan, K., 1988. Flora of Turkey and the East Aegean Islands, Vol. 10 (suppl. 1), *Edinburg University Press, Edinburgh*.
- Demir, A. (2013). Sürdürülebilir gelişmede yükselen değer; biyolojik çeşitlilik açısından Türkiye değerlendirmesi. *İstanbul Ticaret Üniversitesi Fen Bilimleri Dergisi*, 24, 67-74.
- Döldarslan, M., Gül, E. (2015). Yapraklı - Büyükyayla (Çankırı)'nın Vasküler Bitkiler Florası. *Orman ancılık Dergisi* 11(2), 74-91.
- Döldarslan, M., Gül, E. (2021). Yapraklı- Küçükyayla (Çankırı)'nın Florası. *Anadolu Orman Araştırmaları Dergisi* 7(1), 1-14.
- Edis, S., Aytaş, İ., & Özcan, A. U. (2021). Assesment of soil erosion risk using the ICONA model: M eseli (Çubuk/Ankara) Watershed. *Anatolian Journal of Forest Research*, 7(1), 15-22.
- Edis, S., Tuttu, G., Aytaş, İ., Tuttu, U., & Özcan, A. U. (2022). Analysis of temporal and spatial change in Açıçay (Çankırı) Riparian Zone. *Artvin Coruh University Journal of Forestry Faculty*, 23 (1), 1-10.
- Edis, S., Timur, Ö. B., Tuttu, G., Aytaş, İ., Göl, C., & Özcan, A. U. (2023). Assessing the impact of engineering measures and vegetation restoration on soil erosion: a case study in Osmancık, Türkiye. *Sustainability*, 15(15), 12001.
- Edis, S. 2024. İklim Değişikliğinin Kurak ve Yarı Kurak Alanlardaki Hidrolojik Süreçlere Etkileri, (Ed: Gül, E., Döldarslan, M.). Bidge Yayınları, 210 -244, ISBN:978-625-372-159-6, 402 s., Ankara.
- Ekim, T. 2009. Türkiye'nin Nadir Endemikleri. *Türkiye İş Bankası Kültür Yay. No: 11213, İstanbul*.
- Ekim, T., Koyuncu, M., Vural, M., Duman, H., Aytaç, Z., Adigüzel, N. 2000. Türkiye Bitkileri Kırmızı Kitabı, *Türkiye Tabiatını Koruma Derneği ve Van Yüzüncüyl Üniverstesi, Ankara*.
- Elmarsdottir A., Fjellberg, A., Halldorsson, G., Ingimarsdottir, M., Nielsen, O.K., Nygaard, P., Sigurdsson, B.D. (2008). Effects of afforestation on biodiversity. AFFORNORD. Effects of afforestation on ecosystems, landscape and rural development. *TemaNord*, 562, 37-47.
- Erdoğan, N., Ketenoglu, O., Bingöl, Ü. (2008). Kabalı Dağı (Çerkeş-Çankırı) Florası, Ot Sistematisk Botanik Dergisi, 14(2), 63-82.
- Erik, S., Tarikahya, B. (2004). Türkiye Florası Üzerine, *Kebikeç*, 17, 139-163.

- Erişmiş, M. (2023). Long term drought analysis in the Meriç River Basin according to the standard precipitation evapotranspiration index. International Journal of Geography and Geography Education, (50), 313-328.
- Fırıncıoğlu, H.K., Seefeldt, S.S., Şahin, B. (2007). The effects of long-term grazing exclosures on range plants in the Central Anatolian Region of Turkey. Environmental Management, 39, 326-337.
- Gül, E., Dölderslan, M., Uluğ, K. (2019). Yarı Kurak Ağaçlandırma Alanlarında Çölleşme Eğiliminin Değerlendirilmesi: ÇAKÜ Orman Fakültesi, Prof. Dr. Abdülreşit BROHİ Araştırma ve Uygulama Ormanı Örneği. Bartın Orman Fakültesi Dergisi, 21(2), 506-516.
- Güner, A., Aslan, S., Ekim, T., Vural M., Babaç, M.T., (Eds.) 2012. Türkiye Bitkileri Listesi (Damarlı Bitkiler). *Nezahat Gökyigit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayıncılık, İstanbul*.
- Güner, A., Özhatay, N., Ekim, T., Başer, K.H.C. 2000. Flora of Turkey and the East Aegean Islands, vol. 11 (suppl. 2). *Edinburgh University Press, Edinburgh*.
- Harris, J.G., Harris, M.W. 2001. Plant Identification Terminology, (An Illustrated Glossary), *Springs Lake Publishing, Utah*.
- Kar, A. (2018). Desertification: causes and effects. In: Exploring Natural Hazards. *Chapman and Hall/CRC*, 159-206.
- Lal, R. (2004). Carbon sequestration in dryland ecosystems. Environmental management, 33, 528-544.
- Lopez-Bermudez, F., Romero-Díaz, A., Martínez-Fernandez, J., Martínez-Fernandez, J. (1998). Vegetation and soil erosion under a semi-arid Mediterranean climate: a case study from Murcia (Spain). *Geomorphology*, 24(1), 51-58.
- Maestre, F.T., Quero, J.L., Gotelli, N.J., Escudero, A., Ochoa, V., Delgado-Baquerizo, M., Zaady, E. (2012). Plant species richness and ecosystem multifunctionality in global drylands. *Science*, 335(6065), 214-218.
- Mutlu, H. 2006. Çankırı/Yapraklı Ormanlarının Vasküler Bitkiler Florası, Yüksek Lisans Tezi, Ankara Üniversitesi Fen Bilimleri Enstitüsü.
- Noy-Meir, I. (1973). Desert ecosystems: environment and producers. Annual review of ecology and systematics, 25-51.
- Pehlivan, G. 2007. Ilgaz Dağı Milli Parkı Florası, Yüksek Lisans Tezi, Gazi Üniversitesi Fen Bilimi Enstitüsü.

- Sağiroğlu, M., Duman, H. (2004). Karlık Tepesi ve Çevresinin (Çankırı) Florası, Ot Sistematisk Botanik Dergisi, 11(2), 95-122.
- Serkendiz, H., Tatlı, H. (2023). Investigation of Drought in Turkey with the Resilience Index Developed Based on Socio-Economic and Biophysical Components. Resilience, 7(1), 73-91.
- Tekeli, İ., Güler, Ç., Yerli, S.V., Algan, N., Vaizoğlu, S.A., Dündar Kaya, A., Öztürk, B., Mutlu, B., Demirayak, F. 2006. Dünya'da ve Türkiye'de biyolojik çeşitliliği koruma. Türkiye Bilimler Akademisi, TÜBA Raporları No:13, Ankara.
- Thornthwaite, C.W. 1948. *An approach toward a rational classification of climate*. 38(1), 55-94.
- Tutin, T.G. (Ed) 1964-1993, Flora Europaea, vol. 1-5, Cambridge University Press.
- Tuttu, G. 2024. İklim Değişikliği ve Bitkisel Biyoçeşitlilik. In: Gül, E., Dölderslan, M. (Eds.), İklim Değişikliği ve Etkileri: Bütünsel Bir Yaklaşım. *Bidge Yayınları, Ankara*, 159 -195.
- Tuttu, G., Akkemik, Ü. (2017). Çankırı-Korubaşı Tepe ve Civarındaki Jipsli Alanların Florası. Ot Sistematisk Botanik Dergisi, 24 (1), 45-88.
- Tuttu, G., Ursavaş, S. (2022). Çankırı Karatekin Üniversitesi, Orman Fakültesi Araştırma ve Uygulama Ormanının (Çankırı/Eldivan) Florası. Anadolu Orman Araştırmaları Dergisi, 8 (1), 51-65.
- URL. <http://ww2.bgbm.org/herbarium>
- URL. <http://www.vanherbaryum.yyu.edu.tr>
- URL. <https://wfoplantlist.org>
- URL. IPNI (2024). International Plant Names Index. Published on the Internet <http://www.ipni.org>, The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Herbarium. [Retrieved 11 September 2024]. Erişim Tarihi: 08.10.2024.
- Yaltırık, F., Efe, A. 1996. Otsu Bitkiler Sistemiği Ders Kitabı. *İstanbul Üniversitesi yayınları, İstanbul*.

Table 1. Plant species distributed in the study area (LR(Ic): Lower risk, Least Concern)

Family	Taxon	Phytogeographic Region	Endemism and endangered status
Amaryllidaceae	<i>Chenopodium album</i> L. subsp. <i>album</i> var. <i>album</i>	Unknown - multiregional	-
	<i>Allium atroviolaceum</i> Boiss.	Unknown - multiregional	-
	<i>Allium pseudoflavum</i> Vved.	Irano-Turanian	-
	<i>Bupleurum rotundifolium</i> L.	Unknown - multiregional	-
	<i>Caucalis platycarpos</i> L.	Unknown - multiregional	-
	<i>Daucus carota</i> L.	Unknown - multiregional	-
	<i>Eryngium campestre</i> L. var. <i>virens</i> Link.	Unknown - multiregional	-
Apiaceae	<i>Malabaila secacul</i> (Mill.) Boiss.	Unknown - multiregional	-
	<i>Zosima absinthijolia</i> Link	Unknown - multiregional	-
	<i>Cynanchum acutum</i> L. subsp. <i>acutum</i>	Unknown - multiregional	-
	<i>Muscari armeniacum</i> H.J.Weitch	Unknown - multiregional	-
	<i>Ornithogalum oligophyllum</i> E.D.Clarke	Unknown - multiregional	-
	<i>Achillea arabica</i> Kotschy.	Irano-Turanian	-
	<i>Carduus pycnocephalus</i> L. subsp. <i>albidus</i> (M.Bieb.)	Unknown - multiregional	-
Apocynaceae	<i>Carthamus lanatus</i> L.	Unknown - multiregional	-
	<i>Centaurea solstitialis</i> L. subsp. <i>sostitialis</i>	Unknown - multiregional	-
	<i>Centaurea urvillei</i> DC. subsp. <i>armata</i> Wagenitz	East Mediterranean	-
	<i>Centaurea virgata</i> Lam.	Irano-Turanian	-
	<i>Chondrilla juncea</i> L.	Unknown - multiregional	-
	<i>Cichorium intybus</i> L.	Unknown - multiregional	-
	<i>Cirsium arvense</i> (L.) Scop. subsp. <i>vestitum</i> (Wimmer & Grab.) Petrak	Unknown - multiregional	-
Asteraceae	<i>Cirsium pseudocreticum</i> (P.H.Davis & Parris) Yıldız, Dirmenci & Arabacı	Irano-Turanian	Endemic LR (Ic)
	<i>Crepis foetida</i> L. subsp. <i>rhoeadifolia</i> (M. Bieb.) Čelak.	Unknown - multiregional	-
	<i>Crepis micrantha</i> Czerep.	Unknown - multiregional	-
	<i>Crepis pulchra</i> L. subsp. <i>pulchra</i>	Unknown - multiregional	-
	<i>Crupina crupinastrum</i> (Moris) Vis.	Unknown - multiregional	-
	<i>Echinops viscosus</i> DC. subsp. <i>bithynicus</i> (Boiss.) Rech.f.	Unknown - multiregional	-
	<i>Jurinea pontica</i> Hausskn. & Freyn ex Hausskn.	Irano-Turanian	Endemic LR (Ic)
	<i>Lactuca serriola</i> L.	Euro-Siberian	-
	<i>Leontodon asperimus</i> (Willd.) Endl.	Irano-Turanian	-
	<i>Leontodon biscutellifolius</i> DC.	-	-

Table 1. (Continue)

Family	Taxon	Phytogeographic Region	Endemism and endangered status
Asteraceae	<i>Onopordum tauricum</i> Willd.	Euro-Siberian	-
	<i>Picris strigosa</i> M.Bieb. subsp. <i>strigosa</i>	Irano-Turanian	-
	<i>Pulicaria dysenterica</i> (L.) Bernh.	Unknown - multiregional	-
	<i>Rhaponticum repens</i> (L.) Hidalgo	Irano-Turanian	-
	<i>Senecio leucanthemifolius</i> subsp. <i>vernalis</i> (Waldst. & Kit.) Greuter	Unknown - multiregional	-
	<i>Sonchus asper</i> (L.) Hill subsp. <i>glaucescens</i> (Ord.) Ball	Unknown - multiregional	-
	<i>Taraxacum macroleptum</i> Schischk.	Unknown - multiregional	-
	<i>Tragopogon abbreviatus</i> (Boiss.) Coşkunç. & Gültepe	Unknown - multiregional	-
	<i>Tragopogon aureus</i> Boiss.	Unknown - multiregional	Endemic LR (Ic)
	<i>Tragopogon latifolius</i> Boiss. var. <i>angustifolius</i> Boiss.	Irano-Turanian	-
Berberidaceae	<i>Xanthium spinosum</i> L.	Unknown - multiregional	-
	<i>Xeranthemum annuum</i> L.	Unknown - multiregional	-
	<i>Berberis crataegina</i> DC.	Irano-Turanian	-
	<i>Alkanna orientalis</i> Boiss. var. <i>orientalis</i>	Irano-Turanian	-
	<i>Anchusa pusilla</i> Gusul.	Unknown - multiregional	-
	<i>Asperugo procumbens</i> L.	Euro-Siberian	-
	<i>Buglossoides arvensis</i> (L.) I.M.Johnst.	Unknown - multiregional	-
	<i>Cynoglossum montanum</i> L.	Euro-Siberian	-
	<i>Heliotropium dolosum</i> De Not.	Unknown - multiregional	-
	<i>Matiastrium leptophyllum</i> Brand	Irano-Turanian	Endemic LR (Ic)
Boraginaceae	<i>Moltkia coerulea</i> Lehm.	Irano-Turanian	-
	<i>Myosotis lithospermifolia</i> Hornem.	Unknown - multiregional	-
	<i>Nonea caspica</i> G.Don	Irano-Turanian	-
	<i>Onosma isaurica</i> Boiss. & Heldr.	Irano-Turanian	Endemic LR (Ic)
	<i>Aethionema arabicum</i> (L.) Andr. ex DC.	Unknown - multiregional	-
	<i>Alyssum desertorum</i> Stapf	Unknown - multiregional	-
	<i>Alyssum hirsutum</i> M.Bieb.	Unknown - multiregional	-
	<i>Boreava orientalis</i> Jaub. & Spach	Unknown - multiregional	-
	<i>Brassica elongata</i> Ehrh.	Unknown - multiregional	-
	<i>Capsella bursa-pastoris</i> Medik	Unknown - multiregional	-
Brassicaceae	<i>Chorispora tenella</i> DC.	Unknown - multiregional	-
	<i>Crambe tataria</i> Sebeók	Unknown - multiregional	-

Table 1. (Continue)

Family	Taxon	Phytogeographic Region	Endemism and endangered status
	<i>Descurainia sophia</i> (L.) Webb ex Prantl	Unknown - multiregional	-
	<i>Isatis glauca</i> Aucher ex Boiss. subsp. <i>exauriculata</i> (Bornm.) P.H.Davis	Irano-Turmanian	-
	<i>Lepidium draba</i> L.	Unknown - multiregional	-
	<i>Malcolmia africana</i> (L.) W.T.Aiton	Unknown - multiregional	-
	<i>Matthiola longipetala</i> DC. subsp. <i>bicornis</i> Ball	Unknown - multiregional	-
	<i>Noccaea perfoliata</i> (L.) Al-Shehbaz	Unknown - multiregional	-
Brassicaceae	<i>Odontarrhena pateri</i> (Nyár.) Španiel, Al-Shehbaz, D.A.German & Marhold	Irano-Turmanian	Endemic LR (lc)
	<i>Sinapis arvensis</i> L.	Unknown - multiregional	-
	<i>Sisymbrium altissimum</i> L.	Unknown - multiregional	-
	<i>Sisymbrium loeselii</i> L.	Unknown - multiregional	-
	<i>Turritis laxa</i> Hayek	Unknown - multiregional	-
Caprifoliaceae	<i>Lomelesia argentea</i> (L.) Greuter & Burdet	Unknown - multiregional	-
	<i>Lomelesia rotata</i> (M.Bieb.) Greuter & Burdet	Irano-Turmanian	-
	<i>Cerasitum perfoliatum</i> L.	Unknown - multiregional	-
	<i>Dianthus nudiflorus</i> Griff.	Unknown - multiregional	-
	<i>Gypsophila eriocalyx</i> Boiss.	Irano-Turmanian	Endemic LR (lc)
Caryophyllaceae	<i>Minuartia anatolica</i> Woronow var. <i>polymorpha</i> McNeill	Unknown - multiregional	-
	<i>Minuartia hybrida</i> (Vill.) Schischk. subsp. <i>hybrida</i>	Unknown - multiregional	-
	<i>Silene cappadocica</i> Boiss.& Heldr.	Irano-Turmanian	-
	<i>Silene pruinosa</i> Boiss.	Unknown - multiregional	-
	<i>Silene vulgaris</i> (Moench) Gärcke var. <i>vulgaris</i>	Unknown - multiregional	-
	<i>Helianthemum salicifolium</i> (L.) Mill.	Unknown - multiregional	-
	<i>Convolvulus arvensis</i> L.	Unknown - multiregional	-
Cistaceae	<i>Convolvulus holosericeus</i> M.Bieb. subsp. <i>holosericeus</i>	Unknown - multiregional	-
	<i>Convolvulus lineatus</i> L.	Unknown - multiregional	-
Cyperaceae	<i>Scirpoides holoschoenus</i> (L.) Soják	Unknown - multiregional	-
Elaeagnaceae	<i>Elaeagnus angustifolia</i> L.	Unknown - multiregional	-
Equisetaceae	<i>Equisetum palustre</i> L.	Unknown - multiregional	-
Euphorbiaceae	<i>Euphorbia macroclada</i> Boiss.	Irano-Turmanian	-
	<i>Euphorbia myrsinites</i> L.	Unknown - multiregional	-
Fabaceae	<i>Alhagi pseudalhagi</i> (M.Bieb.) Desv. ex Shap.	Irano-Turmanian	-
	<i>Astragalus aduncus</i> Willd.	Irano-Turmanian	-

Table 1. (Continue)

Family	Taxon	Phytogeographic Region	Endemism and endangered status
Fabaceae	<i>Astragalus microcephalus</i> Willd.	Irano-Turanian	-
	<i>Astragalus spruneri</i> Boiss.	Unknown – multiregional	-
	<i>Colutea cilicica</i> Boiss. & Balansa	Unknown – multiregional	-
	<i>Genista sessilifolia</i> DC.	Irano-Turanian	-
	<i>Glycyrrhiza glabra</i> L.	Unknown – multiregional	-
	<i>Hedysarum varium</i> Willd.	Unknown – multiregional	-
	<i>Lotus tenuis</i> Waldst. & Kit. ex Willd.	Unknown – multiregional	-
	<i>Medicago minima</i> (L.) L.	Unknown – multiregional	-
	<i>Medicago sativa</i> L. subsp. <i>sativa</i>	Unknown – multiregional	-
	<i>Melilotus officinalis</i> (L.) Pall.	Unknown – multiregional	-
	<i>Melilotus tauricus</i> (M.Bieb.) Ser.	Unknown – multiregional	-
	<i>Onobrychis hypargyreia</i> Boiss.	Unknown – multiregional	-
	<i>Onobrychis ornata</i> Desv.	Irano-Turanian	Endemic LR (lc)
	<i>Onobrychis oxyodonta</i> Boiss.	Unknown – multiregional	-
Fagaceae	<i>Vicia sativa</i> subsp. <i>nigra</i> Ehrh.	Unknown – multiregional	-
	<i>Quercus robur</i> L. subsp. <i>robur</i>	Euro-Siberian	-
	<i>Centaurium erythraea</i> Rafn subsp. <i>turicum</i> (Velen.) Melderis	Unknown – multiregional	-
Gentianaceae	<i>Erodium cicutarium</i> (L.) L'Hér. subsp. <i>cicutarium</i>	Unknown – multiregional	-
	<i>Hypericum organifolium</i> Willd.	Unknown – multiregional	-
Hypericaceae	<i>Juncus inflexus</i> L.	Unknown - multiregional	-
	<i>Ballota nigra</i> L. subsp. <i>anatolica</i> P.H. Davis	Irano-Turanian	-
Juncaceae	<i>Lamium orientale</i> (Fisch. & C.A.Mey.) E.H.L.Krause	Irano-Turanian	-
	<i>Lamium purpureum</i> L.	Euro-Siberian	-
Lamiaceae	<i>Marrubium parviflorum</i> Fisch. & C.A. Mey. subsp. <i>oligodon</i> (Boiss.) Seybold	Unknown – multiregional	-
	<i>Salvia absconditiflora</i> Greuter & Burdet	Irano-Turanian	Endemic LR (lc)
	<i>Salvia aethiopis</i> L.	Unknown – multiregional	-
	<i>Salvia sclarea</i> L.	Unknown – multiregional	-
	<i>Scutellaria orientalis</i> L. subsp. <i>pinnatifida</i> J.R.Edm.	Unknown - multiregional	-
	<i>Sideritis montana</i> L. subsp. <i>remota</i> (d'Urv.) P.W.Ball	East Mediterranean	-
	<i>Stachys annua</i> (L.) L. subsp. <i>annua</i> (L.) L.	Unknown – multiregional	-
	<i>Teucrium chamaedrys</i> L. subsp. <i>syriacum</i> (K.Koch) Rech.f.	Irano-Turanian	-

Table 1. (Continue)

Family	Taxon	Phytogeographic Region	Endemism and endangered status
Lamiaceae	<i>Teucrium polium</i> L.	Unknown - multiregional	-
	<i>Teucrium scordium</i> L. subsp. <i>scordiooides</i> (Schreb.) Arcang.	Euro-Siberian	-
	<i>Thymus longicaulis</i> C.Presl subsp. <i>chaubardii</i> (Rchb.f.) Jalas	Mediterranean	-
	<i>Ziziphora taurica</i> M.Bieb.	Irano-Turanian	-
	<i>Ziziphora tenuior</i> L.	Irano-Turanian	-
Linaceae	<i>Linum tenuifolium</i> L.	Unknown - multiregional	-
Malvaceae	<i>Alcea apterocarpa</i> Boiss.	Irano-Turanian	-
Nitrariaceae	<i>Peganum harmala</i> L.	Unknown - multiregional	-
Oleaceae	<i>Chrysojasminum fruticans</i> (L.) Banfi	Mediterranean	-
	<i>Fumaria officinalis</i> L.	Unknown - multiregional	-
Papaveraceae	<i>Glaucium grandiflorum</i> subsp. <i>refractum</i> (Nábělek) Mory	Irano-Turanian	-
	<i>Hypecoum procumbens</i> L.	Mediterranean	-
	<i>Papaver laevigatum</i> M.Bieb.	Unknown - multiregional	-
	<i>Papaver rhoeas</i> L.	Unknown - multiregional	-
	<i>Andrachne telephioides</i> L.	Unknown - multiregional	-
Phyllanthaceae	<i>Globularia orientalis</i> L. var. <i>orientalis</i>	Irano-Turanian	-
Plantaginaceae	<i>Plantago lanceolata</i> L.	Unknown - multiregional	-
	<i>Plantago maritima</i> L.	Unknown - multiregional	-
	<i>Veronica hederifolia</i> L.	Unknown - multiregional	-
	<i>Veronica multifida</i> L.	Irano-Turanian	-
	<i>Veronica persica</i> Poir.	Unknown - multiregional	-
	<i>Aegilops speltoides</i> Tausch	Unknown - multiregional	-
	<i>Aegilops triuncialis</i> L. subsp. <i>triuncialis</i>	Unknown - multiregional	-
	<i>Avena fatua</i> L.	Unknown - multiregional	-
Poaceae	<i>Bothriochloa ischaemum</i> (L.) Keng	Unknown - multiregional	-
	<i>Cenchrus orientalis</i> (Rich.) Morrone	Irano-Turanian	-
	<i>Chrysopogon gryllus</i> (L.) Trin.	Unknown - multiregional	-
	<i>Dactylis glomerata</i> L. subsp. <i>hispanica</i> (Roth) Nyman	Unknown - multiregional	-
	<i>Melica ciliata</i> L. subsp. <i>ciliata</i>	Unknown - multiregional	-
	<i>Milium vernale</i> M.Bieb.	Mediterranean	-

Table 1. (Continue)

Family	Taxon	Phytogeographic Region	Endemism and endangered status
Poaceae	<i>Phleum subulatum</i> (Savi) Asch. & Graebn.	Unknown - multiregional	-
	<i>Phragmites australis</i> (Cav.) Trin. ex Steud.	Euro-Siberian	-
	<i>Poa bulbosa</i> L.	Unknown - multiregional	-
	<i>Polypogon monspeliensis</i> (L.) Desf.	Unknown - multiregional	-
	<i>Stipa arabica</i> Trin. & Rupr.	Irano-Turanian	-
	<i>Triticum aestivum</i> L.	Unknown - multiregional	-
	<i>Polygonum pruinosa</i> Boiss. subsp. <i>pruinosa</i>	Unknown - multiregional	-
Polygalaceae	<i>Polygonum bellardii</i> All.	Unknown - multiregional	-
	<i>Rumex dentatus</i> L. subsp. <i>halacsyi</i> (Rech.) Rech.f.	Unknown - multiregional	-
Primulaceae	<i>Androsace maxima</i> L.	Unknown - multiregional	-
Ranunculaceae	<i>Clematis orientalis</i> L.	Unknown - multiregional	-
Resedaceae	<i>Clematis vitalba</i> L.	Unknown - multiregional	-
	<i>Delphinium ajacis</i> L.	Unknown - multiregional	-
	<i>Reseda lutea</i> L. var. <i>lutea</i>	Unknown - multiregional	-
	<i>Paliurus spina-christi</i> Mill.	Unknown - multiregional	-
	<i>Reseda lutea</i> L. var. <i>lutea</i>	Unknown - multiregional	-
	<i>Paliurus spina-christi</i> Mill.	Unknown - multiregional	-
	<i>Potentilla recta</i> L.	Unknown - multiregional	-
Rosaceae	<i>Prunus argentea</i> (Lam.) Rehder	Irano-Turanian	-
	<i>Prunus mahaleb</i> L.	Unknown - multiregional	-
	<i>Rosa canina</i> L.	Unknown - multiregional	-
	<i>Rubus creticus</i> Tourn. ex L.	Unknown - multiregional	-
	<i>Sanguisorba minor</i> Scop. subsp. <i>balearica</i> (Bourg. ex Nyman) Muñoz Garm. & C.Navarro	Unknown - multiregional	-
	<i>Cruciata taurica</i> (Pall. ex Willd.) Ehrend.	Irano-Turanian	-
	<i>Cynanchica bornmuelleri</i> (Velen. ex Bormm.) P.Caputo & Del Giacchio	Irano-Turanian	Endemic LR (c)
Rubiaceae	<i>Cynanchica tenella</i> (Heuff. ex Degen) P.Caputo & Del Giacchio	Unknown - multiregional	-
Salicaceae	<i>Galium incanum</i> Sm. subsp. <i>elatius</i> (Boiss.) Ehrend.	Irano-Turanian	-
	<i>Galium spurium</i> L.	Euro-Siberian	-
Santalaceae	<i>Rubia tinctorum</i> L.	Irano-Turanian	-
Santalaceae	<i>Populus nigra</i> L. subsp. <i>nigra</i>	Unknown - multiregional	-
	<i>Thesium billardieri</i> Boiss.	Irano-Turanian	-

Table 1. (Continue)

Family	Taxon	Phytogeographic Region	Endemism and endangered status
Scrophulariaceae	<i>Scrophularia canina</i> L. subsp. <i>bicolor</i> (Sm.) Greuter	East Mediterranean	-
	<i>Verbascum cheiranthifolium</i> Boiss. var. <i>cheiranthifolium</i>	Unknown - multiregional	-
Tamaricaceae	<i>Tamarix smyrnensis</i> Bunge	Unknown - multiregional	-
Typhaceae	<i>Typha domingensis</i> Pers.	Unknown - multiregional	-
Violaceae	<i>Viola oculta</i> Lehm.	Unknown - multiregional	-
Zygophyllaceae	<i>Tribulus terrestris</i> L. <i>Zygophyllum fabago</i> L.	Unknown - multiregional Irano-Turanian	-



This paper DOI: [10.5281/zenodo.14025324](https://doi.org/10.5281/zenodo.14025324)

Journal New Website web1: <https://ijgsw.eu.org/>
web2: <https://ijgsw.net/>

Journal Old Website: <http://ijgsw.comze.com/> is no longer used
You can submit your paper to email: Jichao@email.com
[Or IJGSW@mail.com](mailto:IJGSW@mail.com)